## Table 13.6g Immunosuppression Use for Maintenance Between Discharge and One Year Following Transplantation, 1997 to 2006 Recipients with Heart-Lung Transplants

	Year of Transplant									
	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Transplants	62	47	51	48	27	33	29	40	35	31
Tx with Follow-up Immunosuppression Info	34	24	28	29	21	21	13	30	26	20
Corticosteroids										
Any in Category	100.0%	95.8%	96.4%	100.0%	95.2%	100.0%	92.3%	100.0%	100.0%	100.09
Steroids	100.0%	95.8%	96.4%	100.0%	95.2%	100.0%	92.3%	100.0%	100.0%	100.09
Cyclosporine										
Any in Category	55.9%	62.5%	53.6%	58.6%	47.6%	52.4%	38.5%	36.7%	50.0%	15.09
Sandimmmune	5.9%	0.0%	0.0%	6.9%	0.0%	0.0%	0.0%	0.0%	0.0%	0.09
Neoral	50.0%	62.5%	53.6%	44.8%	42.9%	47.6%	23.1%	26.7%	30.8%	5.09
Gengraf	0.0%	0.0%	0.0%	6.9%	4.8%	4.8%	15.4%	10.0%	19.2%	10.0º
Eon	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.09
Tacrolimus										
Any in Category	47.1%	37.5%	50.0%	58.6%	76.2%	66.7%	76.9%	73.3%	84.6%	90.09
Tacrolimus	47.1%	37.5%	50.0%	58.6%	76.2%	66.7%	76.9%	73.3%	84.6%	90.09
Antimetabolites										
Any in Category	88.2%	95.8%	89.3%	93.1%	95.2%	95.2%	84.6%	90.0%	100.0%	100.09
Mycophenolate Mofetil	23.5%	54.2%	42.9%	48.3%	47.6%	38.1%	38.5%	43.3%	53.8%	60.09
Mycophenolate Sodium	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.09
Azathioprine	64.7%	41.7%	46.4%	44.8%	47.6%	57.1%	46.2%	46.7%	46.2%	40.09
Cytoxan	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.09
mTOR Inhibitors										
Any in Category	0.0%	4.2%	3.6%	10.3%	0.0%	9.5%	0.0%	6.7%	7.7%	0.09
Sirolimus	0.0%	4.2%	3.6%	10.3%	0.0%	9.5%	0.0%	6.7%	7.7%	0.09
Everolimus	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.09

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages are calculated based on the total number of patients with any immunosuppression use between discharge and one year following transplantation.

Individual drug percentages will not necessarily add up to the drug category percentages, as patients may be prescribed more than one drug within the same category.